

How Torsion May Set Cosmological Constant, and Massive Gravity, with Nod to Strain in GW signals

ANDREW WALCOTT BECKWITH¹

Physics department
Chongqing University
People's Republic of China

Abstract

Based on the idea of cyclic conformal cosmology, we discuss how torsion may allow for a cosmological constant, which links the ideas given by Beckwith and QaZi 2023 to a presentation for Torsion as given by de Sabbata and Sirvaram, Erice 1990 .Our formulation leads to a left over cosmological constant 10^{-121} times vacuum energy . Furthermore, using massive gravity as proportional to the square root of the cosmological constant, we discuss initial gravity formed immediately after Planckian dynamics, as well as considerations as to elementary strain for GW signals initially propagated. At the start of inflation.

Key words, Inflation, Gravitational waves, Penrose CCC

I. Introduction: Review of the purported role of Torsion given by de Sabbata and Sirvaram 1990 Versus a preview of what we will be doing

To begin this look at [1] [2][3] which purports to show a global cancellation of a vacuum energy term, which is akin, as we discuss later to cancelling the following completely [3] [4][5]

$$\rho_{\Lambda} c^2 = \int_0^{E_{\text{Planck}}/c} \frac{4\pi p^2 dp}{(2\pi\hbar)^3} \cdot \left(\frac{1}{2} \cdot \sqrt{p^2 c^2 + m^2 c^4} \right) \approx \frac{(3 \times 10^{19} \text{ GeV})^4}{(2\pi\hbar)^3} \tag{1}$$

$$\xrightarrow{E_{\text{Planck}}/c \rightarrow 10^{-30}} \frac{(2.5 \times 10^{-11} \text{ GeV})^4}{(2\pi\hbar)^3}$$

In [1], the first line is the vacuum energy which is completely cancelled in their formulation of application of Torsion. In our article we are arguing for the second line . In fact, in our formulation our reduction to the second line of Eq. (1) will be to confirm the following change in the Planck energy term given by [1]

$$\frac{\Delta E}{c} = 10^{18} \text{ GeV} - \frac{n_{\text{quantum}}}{2c} ; 10^{-12} \text{ GeV} \tag{2}$$

The term n (quantum) comes from a Corda derived expression as to energy level of relic black holes [4]. We argue that our application of [1] [2] will be commensurate with Eq. (2) which uses the value given in [2] as to the following .i.e. relic black holes will contribute to the generation of a cut off of the energy of the integral given in Eq. (1) whereas what is done in Eq.(1) by [1] [2] is restricted to a different venue which is reproduced below, namely cancellation of the following by Torsion

$$\rho_{\Lambda} c^2 = \int_0^{E_{\text{Planck}}/c} \frac{4\pi p^2 dp}{(2\pi\hbar)^3} \cdot \left(\frac{1}{2} \cdot \sqrt{p^2 c^2 + m^2 c^4} \right) \approx \frac{(3 \times 10^{19} \text{ GeV})^4}{(2\pi\hbar)^3} \quad (3)$$

Furthermore, the claim in [1] is that there is no cosmological constant, i.e. that Torsion always cancelling Eq. (3) which we view is incommensurate with Table 1 as of [3] which is given below . We claim that the influence of Torsion will aid in the decomposition of what is given in Table 1 below from [3] and will furthermore lead to the influx of primordial black holes which we claim is responsible for the behavior of Eq. (2) above

Table 1 from [2] assuming Penrose recycling of the Universe as stated in that document [5]

End of Prior Universe time frame	Mass (black hole) : super massive end of time BH 1.98910 ⁺⁴¹ to about 10 ⁴⁴ grams	Number (black holes) 10 ⁶ to 10 ⁹ of them usually from center of galaxies
Planck era Black hole formation Assuming start of merging of micro black hole pairs	Mass (black hole) 10 ⁻⁵ to 10 ⁻⁴ grams (an order of magnitude of the Planck mass value)	Number (black holes) 10 ⁴⁰ to about 10 ⁴⁵ , assuming that there was not too much destruction of matter-energy from the Pre Planck conditions to Planck conditions
Post Planck era black holes with the possibility of using Eq. (1) to have say 10 ¹⁰ gravitons/second released per black hole	Mass (black hole) 10 grams to say 10 ⁶ grams per black hole	Number (black holes) Due to repeated Black hole pair forming a single black hole multiple time. 10 ²⁰ to at most 10 ²⁵

II. Now for the statement of the Torsion problem as given in [1] with a nod to [6] [7][8], in the massless particle case, initially

The author is very much aware as to quack science as to purported torsion physics presentations and wishes to state that the torsion problem is not linked to anything other than disruption as to the initial configuration of the expansion of the universe and cosmology, more in the spirit of [6], [7] and is nothing else. Hence, in saying this we wish to delve into what was given in [1] with a subsequent follow up and modification: We first follow the description of [1] to remove Torsion physics from the quacks

To do this, note that in [1] the vacuum energy density is stated to be

$$\rho_{vac} = \Lambda_{eff} c^4 / 8\pi G \quad (4)$$

And

$$(da / d\tau)^2 = \left[1 - \left(r_{min}^4 / a^4 \right) \right] \quad (5)$$

With , if S is the so called spin scalar S and identified as the basic h unit of spin

$$r_{min}^4 = 3G^2 S^2 / 8c^4 \quad (6)$$

III. How to modify Eq. (13) in the presence of matter via Yang Mills fields $F_{\mu\nu}^\beta$

If $g = hc$ we have $\beta_1 = r_{min}^2, \beta_2 = r_{min}^4$, and the minimum radius is identified with a Planck Radius so then

$$(da / d\tau)^2 = \left[1 - \left((\beta_1 = l_p^2) / a^2 \right) - \left((\beta_2 = l_p^4) / a^4 \right) \right] \quad (7)$$

Eventually in the case of an unpolarized spinning fluid in the immediate aftermath of the big bang, we would see a Roberson Walker universe given as, if σ is a torsion spin term added due to [1] as

$$\left(\frac{\dot{R}_0}{R_0} \right)^2 = \left(\frac{8\pi G}{3} \right) \cdot \left[\rho - \frac{2\pi G \sigma^2}{3c^4} \right] + \frac{\Lambda c^2}{3} - \frac{\dot{R}_0^2}{R_0^2} \quad (8)$$

IV. What [1] does as to Eq. (8)

In the case of [1] we would see σ be identified as due to torsion so that Eq. (8) reduces to

$$\left(\frac{\dot{R}_0}{R_0} \right)^2 = \left(\frac{8\pi G}{3} \right) \cdot [\rho] - \frac{\dot{R}_0^2}{R_0^2} \quad (9)$$

The claim is made in [1] that this is due to spinning particles which remain invariant so the cosmological vacuum energy, or cosmological constant is always cancelled. Our approach instead will yield

$$\left(\frac{\Lambda_0}{\Lambda_0}\right)^2 = \left(\frac{8\pi G}{3}\right) \cdot [\rho] + \frac{\Lambda_{observed} c^2}{3} - \frac{\hbar c^2}{\Lambda_0} \quad (10)$$

I.e. the observed cosmological constant $\Lambda_{observed}$ is 10^{-122} times smaller than the initial vacuum energy

The main reason for the difference in the Eq. (9) and Eq. (10) is in the following observation. We will go to Table 1 and make the following assertion

Mainly that the reason for the existence of σ^2 is due to the dynamics of spinning black holes in the precursor to the big bang, to the Planckian regime, of space time, whereas in the aftermath of the big bang, we would have a vanishing of the torsion spin term. i.e. the Table 1 dynamics in the aftermath of the Planckian regime of space time would largely eliminate the σ^2 term

V. Filling in the details of the Eq. (9) collapse of the cosmological term, versus the situation given in Eq. (10)

First look at numbers provided by [3] as to inputs, i.e. these are very revealing

$$\Lambda_{pl} c^2 \approx 10^{87} \quad (11)$$

This is the number for the vacuum energy and this enormous value is 10^{122} times larger than the observed cosmological constant. Torsion physics, as given by [3] is solely to remove this giant number. In order to remove it, the reference [3] proceeds to make the following identification, namely

$$\left(\frac{8\pi G}{3}\right) \cdot \left[-\frac{2\pi G \sigma^2}{3c^4}\right] + \frac{\Lambda c^2}{3} = 0 \quad (12)$$

What we are arguing is that instead, one is seeing, instead

$$\left(\frac{8\pi G}{3}\right) \cdot \left[-\frac{2\pi G \sigma^2}{3c^4}\right] + \frac{\Lambda_{pl} c^2}{3} \approx 10^{-122} \times \left(\frac{\Lambda_{pl} c^2}{3}\right) \quad (13)$$

Our timing as to Eq. (12) is to unleash a Planck time interval t about 10^{-43} seconds. As to Eq. (12) versus Eq. (13) the creation of the torsion term is due to a presumed particle density of

$$n_{pl} \approx 10^{98} \text{ cm}^{-3} \quad (14)$$

Finally, we have a spin density term of

$$\sigma_{pl} = n_{pl} h \approx 10^{71} \tag{15}$$

VI. Conclusion 1, what about massive gravity ?

We will assume for the moment that Eq. (12) and Eq. (3) share in common Eq. (14) and Eq. (15) Does this all mesh in with more classical relativity assumptions? First of all, if this is related to the following

If so, by Novello [8][7] we then have a bridge to the cosmological constant as given by

$$m_g = \frac{h \cdot \sqrt{\Lambda}}{c} \tag{16}$$

What we are referring to is in our model what would be in the near the beginning of inflation and we would be trying to reconcile if our use of mini black holes could be made commensurate as to the existence of setting our value of rest massive gravity to about the square root of the Cosmological constant. Bear in mind that if we do so that the present value of the cosmological constant in Eq. (16) is , if we assume Planck units, [5]

$$\begin{aligned} \rho_{\Lambda} c^2 &= \frac{E_{\Lambda}}{V} \approx \frac{1}{2} \cdot \int_0^{E/c} \sqrt{p^2 c^2 + m^2} \cdot \frac{4\pi p dp}{(2\pi h)^3} \approx \frac{E^4}{16\pi^2 \cdot (hc)^3} \\ &\xrightarrow{E=E_p} \frac{(3 \times 10^{27} eV)^4}{(hc)^3} \\ \rho_{\Lambda} c^2 &\approx \frac{(3 \times 10^{27} eV)^4}{(hc)^3} \xrightarrow{\text{Critical-density}} \frac{(2.5 \times 10^{-3} eV)^4}{(hc)^3} \\ E(\text{field - theory}) &\approx m_p \approx 10^{-5} g \approx 10^{27} eV \\ E(\text{vacuum - meas}) &\approx 10^{-35} g \approx 10^{-5} g \approx 10^{-3} eV \\ E(\text{graviton - rest}) &\approx 10^{-65} g \approx 10^{-33} eV \\ \rho_{de} &\equiv 3c^2 m_p^2 L^{-2} \\ L &= 1 / R_b \\ R_b &= a \cdot \int_a^{\infty} \frac{da}{Ha^2} \end{aligned} \tag{17}$$

It appears to be trivial, a mere round off, but I can assure you the difference is anything but trivial. And this is where Table 1 really plays a role in terms of why there is a torsion term to begin with, i.e. will make the following determination, i.e.

The term of ‘spin density’ in Eq. (12) by Eq. (15) is defined to be an ad hoc creation, as to [3]. No description as to its origins is really offered

We state that in the future a task will be to derive in a coherent fashion the following, i.e. the term of

$$\left(\frac{8\pi G}{3}\right) \cdot \left[-\frac{2\pi G\sigma^2}{3c^4}\right] \text{ arising as a result of the dynamics of Table 1, as given in the manuscript}$$

2nd,

The conclusion of [3] states that Eq.(12) would remain invariant for the life of the evolution of the universe. We make no such assumption. We assume that, as will be followed up later that Eq. (13) is due to relic black holes with the suppression of the initially gigantic cosmological vacuum energy,

We state that the term $\left(\frac{8\pi G}{3}\right) \cdot \left[-\frac{2\pi G\sigma^2}{3c^4}\right]$ is due to initial micro black holes, as to the creation of a Cosmological term. This would follow from Eq. (2) being utilized, i.e. what we are seeking is utilization of the following

VI. Dealing with vacuum energy and how would this tie in with Strain of GW

In the case of Pre Planckian space-time the idea is to do the following [9] ,i.e. if we have an inflaton field [10]

$$\left|dp_\alpha dx^\alpha\right| \approx \frac{L}{l} \cdot \frac{h}{c} \cdot \left[\frac{dl}{l}\right]^2$$

$$\xrightarrow{\alpha=0} \left|dp_0 dx^0\right|; \left|\Delta E \Delta t\right| \approx \left(h / a_{mit}^2 \phi(t)\right) (18)$$

$$\Rightarrow \frac{L}{l} \cdot \frac{h}{c} \cdot \left[\frac{dl}{l}\right]^2 \approx \left(h / a_{mit}^2 \phi(t_{mit})\right)$$

Making use of all this leads to [8] to making sense of the quantum number n as given by reference to black holes, [4]

$$E_{Bh} = -\frac{n_{quantum}}{2} \quad (19)$$

3rd

The conclusion of [3] states that Eq.(12) would remain invariant for the life of the evolution of the universe. We make no such assumption. We assume that, as will be followed up later that Eq. (13) is due to relic black holes with the suppression of the initially gigantic cosmological vacuum energy, The details of what follow after this initial period of inflation remain a task to be completed in full generality but we are still assuming as a given the following inputs [1] [10]

Also this would be for a black hole with mass as in Eq. (21) [11]

$$\begin{aligned}
m &\approx \frac{M_P}{\sqrt{N_{\text{gravitons}}}} \\
M_{BH} &\approx \sqrt{N_{\text{gravitons}}} \cdot M_P \\
R_{BH} &\approx \sqrt{N_{\text{gravitons}}} \cdot l_P \\
S_{BH} &\approx k_B \cdot N_{\text{gravitons}} \\
T_{BH} &\approx \frac{T_P}{\sqrt{N_{\text{gravitons}}}}
\end{aligned} \tag{20}$$

Where we may have an increase in mass, for M (black hole) looking like

$$\delta M \approx \frac{6k}{8\pi G_5} \cdot \frac{4\pi r_h^3}{3} \tag{21}$$

That value of mass, assuming a volume of $\frac{4\pi r_h^3}{3}$ has been taken from a brane world, according to [11], with a re scaled gravitational value in 5 dimensions set as G_5 , is allegedly commensurate with the creating of a black hole mass as given in Eq. (20). And in future works to be considered, we will eventually use [11] as to configure how to obtain strain for GW due to the production of relic black holes as well as ideas from [12]. While keeping in mind the observations from [13] from *Grishchuk, L. P.*

Bibliography

[1] de Sabbata, V. and Sirvaram, “Quantum Effects and the Problem of Cosmological Constant”, pp 19-36 in “Gravitation and Modern Cosmology, the Cosmological Constant Problem”: edited by Zichichi, A., de Sabbata, V. and Sacherz, N., Volume 56, Ettore Majorana International Science series (Physical sciences), Plenum Press, New York City New York, USA, 1991

[2] Beckwith, A., “New Conservation Law as to Hubble Parameter, Squared Divided by Time Derivative of Inflaton in Early and Late Universe, Compared with Discussion of HUP in Pre Planckian to Planckian Physics, and Relevance of Fifth Force Analysis to Gravitons and GW”, pp 1 -18 in, *Gravitational Waves - Theory and Observations*, edited by Prof. Carlos Frajuca, published by Intechopen, London UK, 2023 Submitted: October 24th, 2022 Reviewed: October 30th, 2022 Published: December 22nd, 2022 DOI: 10.5992/intechopen.1000577, <https://www.intechopen.com/online-first/1125889>

[3] Beckwith, A. and Ghafoor, Q. (2023) Using Model of a Universe as Similar to a Black Hole, Ask If We Have to Have Singularities, If We Are Looking at Initial Time Step and Entropy, from the Beginning. *Journal of High Energy Physics, Gravitation and Cosmology*, **9**, 708-719. doi: [10.4236/jhepgc.2023.93058](https://doi.org/10.4236/jhepgc.2023.93058).

[3] Cheng Ta-Pei, “Relativity, Gravitation and cosmology, a Basic introduction” Oxford University press united kingdom , 2008

[4] Corda, Christian, “Black Hole spectra from Vaz’s Quantum gravitational collapse”, <https://arxiv.org/abs/2305.02184>, 11 pages, accepted for publication in Fortschritte der Physik - Progress of Physics

[5] Beckwith, A. (2024) How Torsion as Presented by De Sabbata and Sivaram in Erice 1990 Argument as Modified May Permit Cosmological Constant, and Baseline as to Dark Energy. *Journal of High Energy Physics, Gravitation and Cosmology*, **10**, 138-148. doi: [10.4236/jhepgc.2024.101012](https://doi.org/10.4236/jhepgc.2024.101012).

[6] [Sciama, D.W.](#) (1964), "*The physical structure of general relativity*", *Rev. Mod. Phys.*, **36** (1): 463, *Bibcode:1964RvMP...36..463S*, doi:[10.1103/RevModPhys.36.463](https://doi.org/10.1103/RevModPhys.36.463)

[7] de Sabbata, V., And Gasperini, G., Introduction to Gravitation, World Scientific, Singapore, Republic of Singapore, 1989

[8] Novello, M., “The mass of the graviton and the cosmological constant puzzle”, <https://arxiv.org/abs/astro-ph/0504505>

[9] Wesson, Paul, “Five-dimensional Physics: Classical And Quantum Consequences Of Kaluzaklein Cosmology” World Press Scientific, Singapore, Republic of Singapore 2006

[10] Padmanabhan, T., “An Invitation to Astrophysics”, World Press Scientific, World Scientific Series in Astronomy and Astrophysics: Volume 8, Singapore, Republic of Singapore, 2006

[11] Beckwith, Andrew,” How Initial Degrees of Freedom May Contribute to Initial Effective Mass”, <https://vixra.org/abs/2209.0144>

[12] LIGO Scientific and VIRGO Collaborations, “The basic physics of the binary black hole merger GW150914”, *Ann. Phys. (Berlin)* 529, No. 1–2, 1600209 (2017) / DOI 10.1002/andp.201600209 , <https://arxiv.org/ftp/arxiv/papers/1608/1608.01940.pdf>

[13] *Grishchuk, L. P. (1976). "Primordial Gravitons and the Possibility of Their Observation". Sov. Phys. JETP Lett. 23 (6): 293–296. Bibcode:1976ZhPmR..23..326G.* PACS numbers: 04.30. + x, 04.90. + e